

Testing factorization in pA collisions at the LHC

Paloma Quiroga-Arias¹, José Guilherme Milhano^{2,3} and Urs Achim Wiedemann³

¹ Departamento de Física de Partículas and IGFAE, Universidade de Santiago de Compostela 15706 Santiago de Compostela, Spain

² CENTRA, Departamento de Física, Instituto Superior Técnico (IST), Av. Rovisco Pais 1, P-1049-001 Lisboa, Portugal

³ Physics Department, Theory Unit, CERN, CH-1211 Genève 23, Switzerland

Abstract

Global perturbative QCD analyses, based on large data sets from e-p and hadron collider experiments, provide tight constraints on the parton distribution function (PDF) in the proton. The extension of these analyses to nuclear parton distributions (nPDF) has attracted much interest in recent years. nPDFs are needed as benchmarks for the characterization of hot QCD matter in nucleus-nucleus collisions, and attract further interest since they may show novel signatures of non-linear density-dependent QCD evolution. However, it is not known from first principles whether the factorization of long-range phenomena into process-independent parton distribution, which underlies global PDF extractions for the proton, extends to nuclear effects. As a consequence, assessing the reliability of nPDFs for benchmark calculations goes beyond testing the numerical accuracy of their extraction and requires phenomenological tests of the factorization assumption. Here we argue that a proton-nucleus collision programme at the LHC, including a rapidity scan, would provide a set of measurements allowing for unprecedented tests of the factorization assumption underlying global nPDF fits.

Keywords: Collinear factorization, nuclear PDFs, LHC, proton-nucleus collisions

1. The collinear factorization approach to single inclusive particle production and the role of parton distribution functions

The factorized QCD ansatz to proton-proton collisions establishes that the cross section for the production of a hadron in such collisions is given by the convolution of the incoming parton distribution functions with the hard cross section and with the fragmentation function. While it has been confirmed in proton-proton, when it comes to hadron-nucleus collisions collinear factorization is a working assumption yet to be tested. Within such approach, the nuclear dependence of the cross section for hadro-production, typically characterized by the nuclear modification factor R_{pA}^h

$$R_{pA}^h(p_T, y) = \frac{d\sigma^{pA \rightarrow h+X}}{dp_T^2 dy} \bigg/ N_{\text{coll}}^{pA} \frac{d\sigma^{pp \rightarrow h+X}}{dp_T^2 dy}, \quad (1)$$

resides solely in the nuclear modified parton distribution functions (nPDFs) by construction (see eq.(1) in [1]).

Email address: pquiroga@fpaxp1.usc.es¹ ()

Parton distribution functions play a central role in the study of high energy collisions involving hadronic projectiles and from the collinear factorization perspective their nuclear modifications are entirely responsible for the deviations from unity of the nuclear modification factor (1). For protons, sets of collinearly factorized universal PDFs are obtained in global pQCD analyses based on data from DIS and DY production, as well as W/Z and jet production at hadron colliders. On the other hand, the understanding of parton distribution functions in nuclei of nucleon number A , $f_{i/A}(x, Q^2)$, is much less mature.

It is customary to characterize nuclear effects in the parton distribution functions by the ratios

$$R_i^A(x, Q^2) \equiv f_{i/A}(x, Q^2) / f_{i/p}(x, Q^2), \quad (2)$$

which show characteristic deviations from unity in global nPDF analysis for all scales of Q^2 tested so far. These effects are typically referred to as nuclear shadowing ($x < 0.01$), anti-shadowing ($0.01 < x < 0.2$), EMC effect ($0.2 < x < 0.7$) and Fermi motion ($x > 0.7$).

Paralleling the determination of proton PDFs, several global QCD analyses of nPDFs have been made within the last decade [2, 3, 4, 5, 6] based, up until recently, solely on fixed-target nuclear DIS and DY data which, compared to the data constraining proton PDFs, are of lower precision and cover a much more limited range of Q^2 and x . Such limitation manifests in nPDFs with huge uncertainties and in a particularly poor gluon distribution function, since this can only be obtained from logarithmic Q^2 -evolution, for which a wide Q^2 -range is mandatory and not available so far.

To improve on the insufficiency of nuclear DIS and DY data, recent global nPDF analyses [3, 2] have included for the first time data from inclusive high- p_T hadron production in hadron-nucleus scattering measured at RHIC [7, 8, 9] under the collinear factorization assumption.

In view of the importance of nPDFs for characterizing benchmark processes in heavy ion collisions, it is thus desirable to look for both stringent phenomenological tests of the working assumption of global nPDF fits and experimental observables that can be used to constrain nPDFs before the construction of DIS experimental facilities as the LHeC [10] or the EIC [11]. To this end we calculate the single inclusive π^0 production at both RHIC and the LHC within the collinear factorization approach, and we argue that a program of hadron-nucleus collisions at the LHC would provide for such tests with unprecedented quality [1, 12]. All the calculations use LO PDFs from CTEQ6L [13] with nuclear modifications EPS09LO [2] and the KKP fragmentation functions [14]. When the collinear factorization ansatz is adopted, the p_T -dependence of R_{pA}^h traces the x -dependence of nPDFs. Although the precise kinematics is complicated by the convolution of the distributions, the qualitative dependence of the Bjorken- x of partons inside the nucleus with p_T of the final hadron and rapidity reads $x \sim p_T e^{-y} / \sqrt{s}$.

2. Testing collinear factorization with pA collisions

Fig. 1 shows the nuclear modification factor $R_{dAu}^{\pi^0}$ for the production of neutral pions in $\sqrt{s_{NN}} = 200$ GeV deuteron-gold collisions at RHIC, calculated within the factorized ansatz at leading order (LO) (also consistent with the NLO-calculation in [2]). The RHIC data [7] in Fig. 1 have been used in constraining the nPDF analysis EPS09 [2] but they were not employed in a closely related nPDF fit [4], which provides an equally satisfactory description of these RHIC data. Therefore, the agreement of data and calculation in Fig. 1 is in support of collinear factorization: the enhancement of $R_{dAu}^{\pi^0}$ in the region around $p_T \simeq 4$ GeV at mid-rapidity tests momentum fractions in the anti-shadowing region.

However, qualitatively different explanations of the $R_{dAu}^{\pi^0}$ measured at RHIC are conceivable. While the above calculation accounted for $R_{dAu}^{\pi^0}(\eta = 0)$ in terms of a nuclear modification of the *longitudinal* parton momentum distribution only, such nuclear effects can be accommodated within a variety of assumptions, namely initial state parton multiple scattering (k_T broadening) [15] and non-linear low- x QCD evolution [16].

Yet, RHIC data being inconclusive about the dynamical explanation of the nuclear modification of the spectrum, we argue that repeating the same measurement at the LHC at a much higher energy and in the much wider available p_T range, will help to disentangle which of the theories is correct, or in which kinematical range each one is valid [1, 12].

When the single inclusive π^0 spectrum is calculated at the TeV scale, the shape of the nuclear modification factor that would be measured at LHC mid-rapidity, Fig. 2, turns out to be qualitatively different from that observed at RHIC: at more than 40 times higher center of mass energy, final state hadrons at the same transverse momentum test $O(40)$ times smaller momentum fractions x_i . Even though a shift of the maximum of $R_{pPb}^{\pi^0}(\eta = 0)$ to such high values of p_T is




Figure 1: Nuclear modification factor (1) at RHIC mid-rapidity. Data points are PHENIX [7]. Solid line represents the EPS09 LO calculation. The dashed lines are the EPS09 uncertainty [2] (inner) plus data normalization uncertainty (outer).




Figure 2: EPS09 LO prediction for the nuclear modification factor of neutral pion single inclusive production in proton-lead collisions at LHC mid-rapidity. All LHC rapidities are in the center-of-mass (shift to lab $\Delta y \sim 0.47$).

a natural consequence in collinear factorization reflecting nuclear modifications of PDFs, it cannot be accounted for in terms of k_T -broadening, since such approaches predict a mild dependence of the spectrum with the center-of-mass energy [15].

Moreover, when the single-inclusive spectrum is calculated for different values of rapidity the results, Fig. 3, show that the rapidity dependence of R_{pPb}^h allows one to scan the main qualitatively different ranges of standard nPDFs in an unprecedented way. The behavior of the nuclear modification factor when saturation is taken into account [16] is rather different: a strong suppression factor is found which is incompatible with any of the existing nuclear parton distribution sets.




Figure 3: Rapidity dependence of $R_{pPb}^{\pi^0}$ (1) for $\sqrt{s_{NN}} = 8.8$ TeV pPb (LHC). The different plots scan the dependence from $y = -4$ (close to Pb projectile rapidity) up to $y = 4$ (close to proton projectile rapidity). Labels indicate whether the nuclear modification originates mainly from the shadowing (S), anti-shadowing (AS) or EMC regime. Vertical lines illustrate the rapidity-dependent p_T range, which can be accessed experimentally with more than $N_{\text{events}} = 1000$ ($= 10$) per GeV-bin within one month of running at nominal luminosity.

Thus, a measurement at the LHC which shows the rapidity dependent behavior in Fig. 2 and 3 would be in strong support of collinear factorization as opposed to initial state multiple scattering and CGC approaches.

3. The LHC as a tool to constrain the gluon nPDF

Despite the perspectives for qualitative tests of collinear factorization, current global analyses of nuclear parton distribution functions come with significant uncertainties as we can see in Fig.12 in [2]. We have compared in Fig. 4 the nuclear modification factor for two nPDF sets, which show marked differences: in contrast to EPS09 the gluon distribution of HKN07 [6] does not show an anti-shadowing peak but turns for $x > 0.2$ from suppression to strong

enhancement at initial scale $Q^2 = 1 \text{ GeV}^2$. Fig. 4 thus illustrates that within the validity of a collinearly factorized approach, LHC data can resolve the qualitative differences between existing nPDF analyses and can improve significantly and within a nominally perturbative regime on our knowledge of nuclear gluon distribution functions.




Figure 4: $R_{pPb}^{\pi^0}$ (1), from two different sets of gluon nuclear PDFs.

Acknowledgments

This work has been supported by MICINN (Spain) under project FPA2008-01177 and FPU grant; Xunta de Galicia (Consellería de Educación) and through grant PGIDIT07PXIB206126PR, the Spanish Consolider- Ingenio 2010 Programme CPAN (CSD2007-00042) (PQA); and Fundação para a Ciência e a Tecnologia (Portugal) under project CERN/FP/109356/2009 (JGM). PQA would like to thank the organizers of the Hard Probes 2010 conference for financial support.

References

- [1] P. Quiroga-Arias, J. G. Milhano, U. A. Wiedemann, Testing nuclear parton distributions with pA collisions at the LHC [arXiv:1002.2537](#).
- [2] K. J. Eskola, H. Paukkunen, C. A. Salgado, EPS09 - a New Generation of NLO and LO Nuclear Parton Distribution Functions, *JHEP* 04 (2009) 065. [arXiv:0902.4154](#), doi:10.1088/1126-6708/2009/04/065.
- [3] K. J. Eskola, H. Paukkunen, C. A. Salgado, An improved global analysis of nuclear parton distribution functions including RHIC data, *JHEP* 07 (2008) 102. [arXiv:0802.0139](#), doi:10.1088/1126-6708/2008/07/102.
- [4] K. J. Eskola, V. J. Kolhinen, C. A. Salgado, The scale dependent nuclear effects in parton distributions for practical applications, *Eur. Phys. J. C* 9 (1999) 61–68. [arXiv:hep-ph/9807297](#), doi:10.1007/s100520050513.
- [5] D. de Florian, R. Sassot, Nuclear parton distributions at next to leading order, *Phys. Rev. D* 69 (2004) 074028. [arXiv:hep-ph/0311227](#), doi:10.1103/PhysRevD.69.074028.
- [6] M. Hirai, S. Kumano, T. H. Nagai, Determination of nuclear parton distribution functions and their uncertainties at next-to-leading order, *Phys. Rev. C* 76 (2007) 065207. [arXiv:0709.3038](#), doi:10.1103/PhysRevC.76.065207.
- [7] S. S. Adler, et al., Centrality dependence of π^0 and eta production at large transverse momentum in $s(NN)^{1/2} = 200$ -GeV d + Au collisions, *Phys. Rev. Lett.* 98 (2007) 172302. [arXiv:nucl-ex/0610036](#), doi:10.1103/PhysRevLett.98.172302.
- [8] J. Adams, et al., Identified hadron spectra at large transverse momentum in p + p and d + Au collisions at $s(NN)^{1/2} = 200$ -GeV, *Phys. Lett. B* 637 (2006) 161–169. [arXiv:nucl-ex/0601033](#), doi:10.1016/j.physletb.2006.04.032.
- [9] I. Arsene, et al., On the evolution of the nuclear modification factors with rapidity and centrality in d + Au collisions at $s(NN)^{1/2} = 200$ -GeV, *Phys. Rev. Lett.* 93 (2004) 242303. [arXiv:nucl-ex/0403005](#), doi:10.1103/PhysRevLett.93.242303.
- [10] M. Klein, et al., Prospects for a Large Hadron Electron Collider (LHeC) at the LHCEPAC’08, 11th European Particle Accelerator Conference, 23–27 June 2008, Genoa, Italy.
- [11] *The Electron Ion Collider: A white paper*, BNL Report BNL-68933-02/07-REV, Eds. A. Deshpande, R. Milner and R. Venugopalan.
- [12] P. Quiroga-Arias, J. G. Milhano, U. A. Wiedemann, Testing collinear factorization and nuclear parton distributions with pA collisions at the LHC [arXiv:1010.1384](#).
- [13] J. Pumplin, A. Belyaev, J. Huston, D. Stump, W. K. Tung, Parton distributions and the strong coupling: CTEQ6AB PDFs, *JHEP* 02 (2006) 032. [arXiv:hep-ph/0512167](#).
- [14] B. A. Kniehl, G. Kramer, B. Potter, Fragmentation functions for pions, kaons, and protons at next-to-leading order, *Nucl. Phys. B* 582 (2000) 514–536. [arXiv:hep-ph/0010289](#), doi:10.1016/S0550-3213(00)00303-5.
- [15] Y. Zhang, G. I. Fai, G. Papp, G. G. Barnafoldi, P. Levai, High p_T pion and kaon production in relativistic nuclear collisions, *Phys. Rev. C* 65 (2002) 034903. [arXiv:hep-ph/0109233](#), doi:10.1103/PhysRevC.65.034903.
- [16] J. L. Albacete, C. Marquet, Single Inclusive Hadron Production at RHIC and the LHC from the Color Glass Condensate, *Phys. Lett. B* 687 (2010) 174–179. [arXiv:1001.1378](#), doi:10.1016/j.physletb.2010.02.073.